126

Cys-Arg-Ser-Thr-Val-Arg-Met-Ser-Ala-Glu-Gln-Val-Gln-Asn-Val-Trp-Val-Arg-Cys TGC-AGA-AGC-ACC-GTG-CGG-ATG-TCT-GCT-GAA-CAG-GTG-CAG-AAC-GTG-TGG-GTT-CGC-TGC SEQ ID No 1: SEQ ID No 2: .

TGF-§ receptor II homology region

(76) FRIQETTCRRESEADPATCDFQRGYHVPVAVCRSTVRMSAEQVQNVWVRCHWSSSSGSSSSEEMFFGDILGSSTS (150)

(151) RNSYLLGLTPDRSRGEPLYEPSREMRRNFPLGNRRYSNPWPRARVNPGFE (200) SEQ ID No 5

(1) MAMKMLVIFVLGMNHWTCTGFPVYDYDPASLKEALSASVAKVNSQSLSPYLFRAFRSSVKRVNALDEDSLTMDLE (75)

cystatin homology region

BMP-2 homology region

leader sequence

cystatin homology region

SEQ ID NO 6	SEQ ID NO 7	
×	0	
S	S	s v
Z	Z	
>	^	-
S	×	İ
Z	A	S U
>	>	
	S	
<b>-</b>	A	C
0	S	တ
>	ب	ပ
	A	
A	щ	
I	$\times$	၁
Z	-	
<u> </u>	S	ပ
S	¥	Ų
z	ے	
	Δ	
Ξ	>	IJ
Ω	_	
A	>-	
	>	U
م	م	
<u>"</u>	14	_
human BMP-2	human BMP-2	

FIG. 2

,								i			ĺ						İ					1	
bovine fetuin	114 C	ပ	۵	D I H V L K Q D	I	>		×	6	۵	ბ ე	ò	FSVLFT	10	1 /				К			SEQIC	SEQ ID NO 8
human TGF- β receptor II	84	ပ	*	>	V A V	>	3	∝	×	Z	۵	ш	Z	7 L I		ш			) \			SEQ ID	SEQ ID NO 9
		_		U		_		U	U	U	တပ	ပ		0,0	s o				_				
human TGF- § receptor II	84	ว	٨	V A V W	^	*	~	KN	Z	Q	*	*	- <del>*</del>	л —	Z	7 1 1			E	ΛL	)	C SEQ ID NO 9	6 ON 0
bovine BBP		၁	8	ST	-	Λ	~	Σ	S	A	п	0	>	N	7	ΜΛ	\ \ \	^	× ×	*		C SEQ ID NO 10	NO 10
		ļ		ပ	s u		_							J J		ပ		رن د	ωv				
							*	0	(	١,												ı	



FIG. 4




FIG. 6





4/7





FIG. 7

FIG. 8A







FIG. 8B

WO 2005/072403 PCT/US2005/002722



WO 2005/072403 PCT/US2005/002722





FIG. 11

IQETTCRRESEADPATCDFQRGYHVPVAVCRSTVRMSAEQV (SEQ ID No 3)
CGEPLYEPSREMRRN" (SEQ ID NO 4 (SEQ ID No 4)



## **SUBSTITUTE SHEET (RULE 26)**

